



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES AND CITIES OF THE UNITED STATES (untabulated).

COLORADO—*Colorado Springs*.—Month of November, 1910. Population, 29,078. Total number of deaths from all causes 46, including tuberculosis 21. Cases reported: Scarlet fever 13, typhoid fever 6.

Month of December, 1910. Total number of deaths from all causes 41, including tuberculosis 17, typhoid fever 1. Cases reported: Scarlet fever 16, typhoid fever 1.

INDIANA.—Month of December, 1910. Population, 2,700,876. Total number of deaths from all causes 3,017, including diphtheria 44, measles 10, scarlet fever 10, tuberculosis 350, typhoid fever 61. Cases reported: Diphtheria 177, smallpox 45, in 7 counties.

KANSAS.—Month of December, 1910. Population, 1,690,949. Total number of deaths from all causes not reported. The deaths from contagious diseases include diphtheria 15, measles 3, scarlet fever 6, smallpox 1, tuberculosis 53, typhoid fever 28. Cases reported: Diphtheria 141, measles 157, scarlet fever 389, smallpox 177, tuberculosis 243, typhoid fever 107.

NEW YORK.—Month of December, 1910. Population, 9,113,614. Total number of deaths from all causes 12,751, including diphtheria 180, measles 47, scarlet fever 93, tuberculosis 1,345, typhoid fever 131. Cases reported: Diphtheria 1,668, measles 2,953, scarlet fever 2,128, smallpox 2, tuberculosis 2,992, typhoid fever 577.

PENNSYLVANIA.—Reports from the State department of health show as follows:

Mortality.—Month of November, 1910. The total number of deaths reported was 9,221, including typhoid fever 204, scarlet fever 87, diphtheria 261, measles 44, whooping cough 78, influenza 74, malarial fever 7, tuberculosis of the lungs 655, tuberculosis of other organs 91, cancer 397, diabetes 61, meningitis 50, acute anterior poliomyelitis 22, pneumonia 1,198, diarrhea and enteritis, under 2 years 284, diarrhea and enteritis, over 2 years 74, Bright's disease 530, early infancy 577, suicide 76, accidents in mines and quarries 73, railway injuries 100, other forms of violence 450, all other diseases 3,822.

Morbidity.—Month of December, 1910. Cases of communicable diseases reported, 10,443, distributed as follows: Anterior poliomyelitis 15, anthrax 2, cerebrospinal meningitis 10, chickenpox 1,028, diphtheria 1,535, erysipelas 154, German measles 58, malarial fever 5, measles 2,275, mumps 343, pneumonia 868, puerperal fever 6, rabies 1, scarlet fever 1,096, smallpox 2, tetanus 4, trachoma 155, trichiniasis 3, tuberculosis 1,079, typhoid fever 1,010, whooping cough 794.